

Do Hardy-type proposals test nonlocality of single-particle?

Antoine Suarez
 Center for Quantum Philosophy,
 CH-Berninastrasse 85, 8057 Zurich, Switzerland
 suarez@leman.ch, www.quantumphil.org
 (Dated: April 12, 2012)

It is argued that Hardy-type proposals assume either nonlocality at detection, and then beg the question, or “local empty waves”, and then have to accept “many worlds” and cannot prove nonlocality.

An experiment demonstrating single-photon nonlocality at detection has been recently presented and discussed [1, 2]. Against the claim of demonstrating single-particle nonlocality in a so simple way one could object that a “more sophisticated” probe of single-particle nonlocality has been proposed by Lucien Hardy in 1994 [3], and improved recently by other authors [4, 5].

However, Hardy’s argument is based on the following axiom in the experimental context of Figure 1:

“Detecting a particle on u_1 and detecting a particle on u_2 never happens” because “no more than one photon is emitted from the source at any time.” [3, 4]

This axiom takes clearly for granted the result of our experiment [1, 2], and therefore Hardy’s argument either tacitly assumes as an axiom the nonlocality it aims to prove, or it assumes “empty waves” as local hidden variables. Since the argument leads to a contradiction, this means that at the very most Hardy’s proposal rules out local hidden variables, although without inequalities.

Hardy’s argument is conceived as a ladder of four gedanken-experiments using the scheme sketched in Figure 1 [3, 4]. Alice and Bob each have two choices, which lead to the four possible *Experiments 1-4* described below. The input state incident on s , and the corresponding coherent states incident on a and b are chosen in a way that quantum mechanics yields the outcome predictions correspondingly listed to the following *Experiments 1-4*:

Experiment 1: Alice and Bob both decide to set a detector on their paths u_1 respectively u_2 (dashed detectors).

Quantum mechanical prediction: If Alice detects a particle on u_1 , then she can infer that Bob detects nothing on u_2 , and viceversa, if Bob detects a particle on u_2 , he can infer that Alice detects nothing on u_1 . (Notice that if Alice (Bob) detects nothing on u_1 (u_2), she (he) cannot infer that Bob (Alice) detects a particle on u_2 (u_1)).

Regarding this *Experiment 1* Hardy’s argument states: “In this case, it is clear that they cannot both detect a photon since no more than one photon is emitted from the source at any time. This means that detecting a particle on u_1 and detecting a particle on u_2 never happens.” [3, 4]




FIG. 1: Hardy’s scheme: Alice/Bob can either directly set a detector on her/his path (dashed detectors), or she/he can make a homodyne detection by combining her/his path with a local oscillator at a 50:50 beam splitter BS1 respectively BS2 (solid detectors). The source (not represented) is supposed to be single-photon. The vacuum is incident on the mode t ; the input state is incident on mode s with no more than one photon at any time; the coherent states incident on modes a and b act as local oscillators.

Experiment 2: Alice makes a homodyne detection at c_1 and d_1 , by combining the state on path u_1 with a local oscillator (coherent state) a , at her 50:50 beam splitter BS1. Bob, meanwhile, makes the same measurement as in *Experiment 1*.

Quantum mechanical prediction: If Alice detects a particle at d_1 and nothing at c_1 , then Bob must detect one particle.

Experiment 3: The roles of Alice and Bob are reversed: Alice sets her detector on path u_1 and Bob makes a homodyne detection at c_2 and d_2 .

Quantum mechanical prediction: if Bob detects one particle at d_2 and nothing at c_2 , then he can infer that Alice must have detected a particle in path u_1 .

Experiment 4: Both Alice and Bob choose to make homodyne detections.

Quantum mechanical prediction: According to quan-

tum mechanics the experiment can have different outcomes. One of the possible ones is that Alice records one particle at d_1 and nothing at c_1 , while Bob records one particle at d_2 and nothing at c_2 .

Suppose now the following reasoning: The result of *Experiment 4* contradicts *Experiment 1*: Indeed, according to *Experiment 2* Alice infers from her measurement that a single particle must have traveled along path u_2 towards Bob. At least, this is true in the sense that, had Bob put his detector in path u_2 , he would have been guaranteed to detect a particle. However, according to *Experiment 3*, at the same time Bob infers from his measurement that a single particle must have traveled along path u_1 towards Alice. According to *Experiment 1* Alice and Bob cannot both be right.

Hardy pointed out that this reasoning does tacitly assume locality, and without this assumption the contradiction disappears: Alice might deduce from her result that, had Bob measured the number of particles in path u_2 , he would definitely have detected exactly one. However, if Bob had measured u_2 instead of the homodyne measurement he did make, there might have been a nonlocal influence from Bob's end to Alice's end and then she might have obtained a different measurement outcome.[3, 4]

The first step of Hardy's ladder, i.e., *Experiment 1*, has clearly the character of an axiom stating the principle of "one photon one count", i.e., conservation of energy in each individual quantum event, in the very same experiment presented in [2]. This could mean that Hardy's axiom *Experiment 1* tacitly assumes as an axiom the nonlocality at detection demonstrated in [1]. But then Hardy's argument would assume just the nonlocality it searches to prove, and not be able of falsifying *local theories without hidden variables* even in principle. Notice that this is not the case of Bell-type experiments, as discussed in [2].

Consequently to avoid begging the question Hardy-type experiments should assume "local empty waves" (i.e., local hidden variables). Consider the statement at the core of Hardy's argument:

"if Bob had measured u_2 instead of the homodyne measurement he did make, there might have been a nonlocal influence from Bob's end to Alice's end and then she might have obtained a different measurement outcome".

This amounts to explain things as follows in terms of "local empty waves":

Suppose the empty wave travels along path u_1 and the particle along u_2 . If these paths are not monitored, nonlocal coordination between the decisions at the beam-splitters BS1 and BS2 is possible and the result expected in *Experiment 4* may appear. By contrast if Alice had put her detector on u_1 , the "empty wave" traveling this

path would not have reached BS1 and therefore the non-local coordination between the decisions at BS1 and BS2 would disappear. Consequently Bob would obtain a different result, that is one that does not allow him to conclude there is a particle on u_1 .

In other words, what Hardy-type proposals test are *local hidden variables* models deriving from the assumption of outcome's decision at the beam-splitters, although a Hardy-type experiment would refute local hidden variables not through an statistical result (like the conventional Bell-type experiments do), but through a single measurement like the GHZ argument does [6].

Nonetheless, (as discussed in [2]) *local hidden variables* lead to "many worlds", and accepting "many worlds" means to accept locality. Therefore arguments refuting only *local hidden variables* but not "many worlds" are not useful to prove nonlocality. We repeat the essential of this argument:

Regarding "many worlds" John Bell said:

"The 'many world interpretation' seems to me an extravagant, and above all an extravagantly vague, hypothesis. I could also dismiss it as silly. And yet... It may have something distinctive to say in connection with the 'Einstein Podolsky Rosen puzzle', and it would be worthwhile, I think, to formulate some precise version of it to see if this is really so." ([7] p. 194).

As it appears, Gilles Brassard and Paul Raymond-Robichaud have now achieved to formulate such a "precise version" with their theory of "parallel lives" [8]. This version has really something distinctive to say in connection with the "EPR puzzle", that is: Refutation of local hidden variables (by the violation of Bell's inequalities or other means) doesn't mean refutation of locality.

Hardy-type proposals refute *local hidden variables* ("empty waves") but not "many worlds" or "parallel lives". Hence they cannot be considered to prove nonlocality.

Whether Hardy-type experiments can be considered "single-particle" ones is matter of debate [4, 9] (likely because the concept of "single-particle experiment" has not been well defined). As a matter of fact Hardy-type experiments use 4 detectors, like the Bell-type experiments (instead of only 2, like the experiment presented in [1, 2]). But in any case this issue is not of interest for the sake of our discussion.

In conclusion, Hardy-type experiments either assume the nonlocality they aim to prove, or assume "local empty waves" and don't prove nonlocality. By contrast "[the] nonlocality of a single particle expressed by Einstein at the 1927 Solvay Conference" [3, 4] is demonstrated by the experiment presented in [1]. The nonlocality this experiment proves [2] is more basic than Bell's nonlocality and rules the whole quantum mechanics: It appears already at the level of single particle experiments (requiring only 2 detectors), and not first in entanglement experiments involving 2 or more particles (requiring at least 4 detec-

tors). This may open the road to deriving the “abstract quantum algebra” from principles with a clear physical content like free will, conservation of energy, and locality emergent from nonlocality [10].

Acknowledgments: I am thankful to Thiago Guerreiro, Bruno Sanguinetti, Nicolas Sangouard, and Hugo Zbinden for stimulating discussions.

-
- [1] T. Guerreiro, B. Sanguinetti, H. Zbinden, N. Gisin and A. Suarez, Single-photon space-like antibunching. arXiv:1204.1712v1 (2012)
 - [2] A. Suarez, ‘Empty waves’, “many worlds”, “parallel lives”, and nonlocal decision at detection. To appear in: A. Suarez and P. Adams (Eds.) *Is Science compatible with free will? Exploring free will and consciousness in light of quantum physics and neuroscience*. Springer, New York, 2012, Chapter 5. arXiv:1204.1732v1 (2012).
 - [3] L. Hardy, Nonlocality of a single photon revisited, *Phys. Rev. Let.* 73,(1994) 2280
 - [4] J. Dunningham and V. Vedral, Nonlocality of a single particle, *Phys. Rev. Let.* 99, (2007) 180404-2
 - [5] L. Heaney, A. Cabello, M. França Santos, and V. Vedral, Extreme nonlocality with one photon, arXiv:0911.0770v3 (2011).
 - [6] D. Greenberger, M. Horne, A. Zeilinger and M. Zukowski, A Bell Theorem Without Inequalities for Two Particles, Using Inefficient Detectors, arXiv:quant-ph/0510207v2 (2008).
 - [7] J.S. Bell, *Speakable and unspeakable in quantum mechanics*, Cambridge: University Press, 1987.
 - [8] G. Brassard and P. Raymond-Robichaud, Can Free Will Emerge from Determinism in Quantum Theory? To appear in: A. Suarez and P. Adams (Eds.) *Is Science compatible with free will? Exploring free will and consciousness in light of quantum physics and neuroscience*. Springer, New York, 2012, Chapter 4. arXiv:1204.2128v1.
 - [9] D. Greenberger, M. Horne, A. Zeilinger, Nonlocality of a single photon? *Phys. Rev. Let.* 75, (1995) 2064
 - [10] A. Suarez, Deriving Bells nonlocality from nonlocality at detection. arXiv:1009.0698v1 (2010).